

New Physics In Top at the Tevatron

Andrew Ivanov
Kansas State University
(for the CDF and D0 Collaborations)

Aspen Winter Conference
January 19, 2010


Searches for New Physics in the Top Quark Sector




**Intrinsic Properties
of Top Quark**

**New Production
Mechanism**


**Non-SM
Top Decays**



Z', KK,... ?
Resonant Production
Forward-Backward Assymetry



**Charge, Mass,
Width, Lifetime ...**



t -> H⁺b - ?

H⁺ -> cs, $\tau\nu$, WA

**New Physics Admixture
in a Sample of Top Quarks**




**T, t', stop, ...
SUSY with (di-)lepton + jets
+ met signatures**

“The pure and
simple truth is
rarely pure and
never simple”
- Oscar Wilde




New Production Mechanism

Standard Model



- Resonant $t\bar{t}$ production is expected from the decays of massive Z -like bosons
- Topcolor
Hill, Parke PRD 49, 4454 (1994)
- KK gluon excitation in RS model
Lillie, Randal, Wang JHEP 0709,074 (2007)




Search in lepton+jets events

Event Selection:



- Lepton (e or μ) $p_T > 25$ GeV, $|\eta| < 1.1(2.0)$
- 3 or more jets with $E_t > 20$ GeV, $|\eta| < 2.5$
- Leading jet $E_t > 40$ GeV
- At least one b-tagged (NN) jet
- Missing $E_t > 20$ (25 GeV) and not collinear to lepton direction
- Solve for p_z of the neutrino and construct $t\bar{t}$ invariant mass from four-momenta

$M_{Z'} > 820$ GeV at 95%CL
 $\Gamma_{Z'} = 1.2\% M_{Z'}$

= 3 jets





≥ 4 jets






Search in all-hadronic mode

- Multi-jet trigger
- Event Selection:
 - 6 or 7 jets with $E_t > 15 \text{ GeV}$, $|\eta| < 2.0$
- Build the tag rate matrix using the QCD enriched data sample and apply it to pre-tag events
- Employ the ANN machinery to improve S/B ratio
- Reconstruct $t\bar{t}$ mass using matrix element approach



Forward-Backward Assymetry

- The SM prediction:
 - Symmetric at LO
- At NLO in $t\bar{t}$ rest frame
- Asymmetry is due to interference of ME amplitudes for the same final state
- $A_{fb} = 0.05 \pm 0.015$
Kuhn, Pordrigo PRL 81,89 (1998)
- Several BSM production mechanisms predict an observable asymmetry (Z' , Axigluons)
- Especially interesting for wide resonances





$$A_{fb} = \frac{N_t(p) - N_t(\bar{p})}{N_t(p) + N_t(\bar{p})}$$



Forward-Backward Asymmetry



$$A_{fb} = \frac{N_t(p) - N_t(\bar{p})}{N_t(p) + N_t(\bar{p})}$$

- Use lepton+jets channel
- Lepton charge provides an information about top/anti-top
- Kinematic event reconstruction similar to top mass analyses
- Use the system of jets from hadronically decaying top quark





Forward-Backward Asymmetry




- Employ an unfolding procedure to go from reconstructed to parton level:
 - 1. Subtract the background
 - 2. Unfold for bin-to-bin migration effects due to detector resolution and event selection effects

$$A_{fb} = 19.3\% \pm 6.5\% \text{ (stat)} \pm 2.4\% \text{ (syst)}$$

- Systematic uncertainty includes various assumptions about the shape of $-Q_l y_{had}$ distribution

Top Charge

Standard Model




- $t \rightarrow bW$ is conceivable due to ambiguity in assigning b-jets to W
 - That corresponds to an exotic (XM) quark of charge $-4/3$
 - While the SM top quark could have even higher mass
 - Motivated by the improvement in EWK fit due to modified b-couplings
-
- **Chang et al, PRD 59, 091503 (1999)**
 - **Ma et al, hep-ph/9909537**




Top Charge Measurement

- Event Selection:
lepton+ ≥ 4 jets
 - ≥ 1 SECVTX b-jet
 - and ≥ 1 SLT b-jet
- Charge Reconstruction:
 - Identifying leptonic and hadronic b (kinematic fitter based on invariant mass constraints)
 - Determining flavor (b or anti-b)



SECOndary VerTeX tagger (SECVTX)
 B-Jet Tag Efficiency 50%
 MiTag Rate (QCD jets) 2.5%



SLT (Soft Lepton Tagger)
 B-Jet Tag Efficiency 40% (e) , 73% (μ)
 MisTag Rate 0.3 % (e), 0.4% (μ)





Top Charge Measurement

- Asymmetry measurement
- Figure of merit is ϵD^2 ,
- Where ϵ ($\sim 2\%$) is the reconstruction efficiency
- $D = 2P - 1$ is the dilution, P ($\sim 60\%$) is a purity
- D is calibrated in data di-jet events

$$A_t = \frac{1}{D} \frac{N_{SM} - N_{XM} - Bkg \times D_{Bkg}}{N_{SM} + N_{XM} - Bkg} \quad (3)$$



- $N_{SM} = 29$
- $N_{XM} = 16$

New





Non-SM Top Decays

Search for Charged Higgs





- Analyze event yields in several channels simultaneously
- Data prefers no H^+ Contribution






Non-SM Top Decays


Search for Charged Higgs



PLB 682, 278 (2009)




Non-SM Top Decays (nMSSM Higgs)



- Light Higgs (A with mass $m_A < 2m_b$) is motivated by inconsistencies in Higgs mass fit
- $H \rightarrow AA$ with $A \rightarrow \tau\tau$ could sufficiently change $Br(H \rightarrow bb)$ to avoid SM LEP limits


Dermisek & Gunion, PRL 95, 041801 (2005), PRD 73, 111701 (2006)

- 2.8 σ excess in $Br(W^\pm \rightarrow \tau \nu)$ at LEP using WW events
- Can be re-conciled with $H^\pm \rightarrow \tau \nu$
 - Dermisek, arXiv:hep-ph/0807.2135
- At tree level $m_H = \sqrt{m_W^2 + m_A^2}$
- Light A requires $m_H \approx m_W$ and opens decay $H^\pm \rightarrow W^* A$
- For low $\tan\beta$ $Br(t \rightarrow H^\pm b)$ could be as high as 40%






Search for nMSSM Higgs in Top Decays





- Standard lepton+jets selection
- Plus at least one isolated soft track with p_T in $[3,20]$ GeV



- In SM additional tracks come mostly from soft parton interactions - Underlying Event
- Employ invariance of track P_t spectrum



Search for nMSSM Higgs in Top Decays





- The performance is validated in various control regions
- The UE contribution is floating freely in the fit
- Validate by measuring consistent Z/gamma cross section



Search for nMSSM Higgs in Top Decays

New




- Signal Region
- Lepton + ≥ 3 jets
- ≥ 1 b-jet
- No evidence for nMSSM Higgs in top decays
- First limits on BR($t \rightarrow H^\pm b$) in this decay channel





Search for t-prime


- Signal Region:
- Lepton ≥ 4 jets
- Use H_T – scalar sum of all transverse momenta
- Perform mass reconstruction similar to top mass
- Excess is larger than due to strong production mechanism





Dobrescu et al, JHEP 0906, 001 (2009)



Search for $t't$




- Lepton + ≥ 4 jets channel
- Search for extra b-jets from Higgs
- Perform simultaneous fit to H_T in $=4$, ≥ 5 jets and $=1$ -tag, $=2$ -tag and ≥ 3 -tag regions




Conclusion

- Our knowledge of the top quark has come a long way since its discovery in 1995
- Many top quark properties are measured to much higher precision
- No evidence for new physics, although there are some hints (Afb asymmetry, t')
- With much higher $t\bar{t}$ cross section lots of LHC new physics searches producing (di-)lepton + jets signatures will be searches in the top sample
- The Tevatron experience will be invaluable for establishing discoveries at LHC





Thank You!



Backup Slides



Asymmetry Dependence on $M_{t\bar{t}}$




- Idea:
 - Study observed A_{fb} asymmetry as a function of $M_{t\bar{t}}$ mass
- Split events into four bins:
 - Forward, low $M_{t\bar{t}}$
 - Forward, high $M_{t\bar{t}}$
 - Backward, low $M_{t\bar{t}}$
 - Backward, high $M_{t\bar{t}}$




Assymetry Dependence on $M_{tt\bar{b}ar}$

Rapidity for Tagged Events with $M_{tt} < 450$ GeV



$$A_{fb} = 15\% \pm 10\% \text{ (stat)} + 6\% \text{ (syst)}$$


Rapidity for Tagged Events with $M_{tt} > 450$ GeV



$$A_{fb} = 24\% \pm 12\% \text{ (stat)} + 8\% \text{ (syst)}$$



- Unfolding is done in four bins similar to A_{fb} measurement

Asymmetry in low vs high M_{tt} for $M_{tt} = 450$ GeV





Asymmetry Dependence on $M_{t\bar{t}}$



• $M_{t\bar{t}}$ Scan Results